arXiv:1606.04673v1 [math.NT] 15 Jun 2016

SYMMETRIC IDENTITIES FOR AN ANALOGUE OF CATALAN POLYNOMIALS

TAEKYUN KIM, DAE SAN KIM, AND JONG-JIN SEO

Abstract. In this paper, we consider an analogue of Catalan polynomials and give some identities of symmetry for those polynomials by using fermionic p-adic integrals on the ring of p-adic integers.

1. Introduction

Let p be an odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will denote the ring of p-adic integers, the field of p-adic rational numbers, and the completion of the algebraic closure of $\mathbb{Q}_p$. The p-adic norm $|\cdot|_p$ is normalized as $|p|_p = \frac{1}{p}$. Let f be a continuous function on $\mathbb{Z}_p$. Then the fermionic p-adic integral of f on $\mathbb{Z}_p$ is defined by Kim

$$\begin{aligned}\int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) &= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\, \mu_{-1}(x) \\ &= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\,(-1)^x, \quad (\text{see [6–10, 13, 14]}).\end{aligned} \tag{1.1}$$

From (1.1), we note that

$$\int_{\mathbb{Z}_p} f_1(x)\, d\mu_{-1}(x) + \int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) = 2f(0), \tag{1.2}$$

where $f_1(x) = f(x+1)$, (see [6–10, 13, 14]).

By (1.2), we get

$$\int_{\mathbb{Z}_p} f(x+n)\, d\mu_{-1}(x) + (-1)^{n-1} \int_{\mathbb{Z}_p} f(x)\, d\mu_{-1}(x) = 2\sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \tag{1.3}$$

where $n \in \mathbb{N}$, (see [6–10, 13, 14]).

As is well known, the Catalan numbers are defined by the generating function to be

$$\frac{1-\sqrt{1-4t}}{2t} = \frac{2}{1+\sqrt{1-4t}} = \sum_{n=0}^{\infty} C_n t^n, (\text{see [1–5, 11, 12, 15, 16]}), \tag{1.4}$$

where $C_n = \binom{2n}{n}\frac{1}{n+1}$, $(n \geq 0)$.

In addition, the Catalan polynomials are given by the generating function

$$\frac{2}{1+\sqrt{1-4t}}(1-4t)^{\frac{x}{2}} = \sum_{n=0}^{\infty} C_n(x)\, t^n, (\text{see [1–5, 11, 12, 15, 16]}). \tag{1.5}$$

Thus, by (1.4) and (1.5), we get

$$C_n(x)=\sum_{m=0}^{n}\sum_{j=0}^{m}\left(\frac{x}{2}\right)^j S_1(m,j)(-4)^m\frac{C_{n-m}}{m!}, \tag{1.6}$$

where $S_1(m,n)$ is the Stirling number of the first kind.

For $w\in\mathbb{N}$, we define w-Catalan polynomials which are given by the generating function

$$\frac{2}{1+(1-4t)^{\frac{w}{2}}}(1-4t)^{\frac{w}{2}x}=\sum_{n=0}^{\infty}C_{n,w}(x)t^n. \tag{1.7}$$

From (1.7), we note that $C_{n,1}(x)=C_n(x)$.

When $x=0$, $C_{n,w}=C_{n,w}(0)$ are called w-Catalan numbers.

From (1.2), we note that

$$\begin{aligned}\int_{\mathbb{Z}_p}(1-4t)^{\frac{w}{2}(x+y)}d\mu_{-1}(y)&=\frac{2}{(1-4t)^{\frac{w}{2}}+1}(1-4t)^{\frac{w}{2}x}\\&=\sum_{n=0}^{\infty}C_{n,w}(x)t^n,\end{aligned} \tag{1.8}$$

where $t\in\mathbb{C}_p$ with $|t|_p<p^{-\frac{1}{p-1}}$.

In particular, for $w=1$, we have

$$\begin{aligned}\int_{\mathbb{Z}_p}(1-4t)^{\frac{x+y}{2}}d\mu_{-1}(y)&=\frac{2}{1+(1-4t)^{\frac{1}{2}}}(1-4t)^{\frac{x}{2}}\\&=\sum_{n=0}^{\infty}C_n(x)t^n.\end{aligned} \tag{1.9}$$

For $k\geq 0$, $d,w\in\mathbb{N}$ with $d\equiv 1 \pmod 2$, $w\equiv 1 \pmod 2$, we define the function $S_{k,d}(w-1)$ as follows:

$$S_{k,d}(w-1)=\sum_{i=0}^{w-1}\binom{\frac{di}{2}}{k}(-1)^i. \tag{1.10}$$

When $d=1$,

$$S_{k,1}(w-1)=S_k(w-1)=\sum_{i=0}^{w-1}\binom{\frac{i}{2}}{k}(-1)^i.$$

In this paper, we give some identities of symmetry for the w-Catalan polynomials which are derived from the fermionic p-adic integrals on $\mathbb{Z}_p$.

2. Identities of symmetry for the w-Catalan polynomials

From (1.9), we note that

$$\int_{\mathbb{Z}_p}\binom{\frac{x+y}{2}}{n}d\mu_{-1}(y)=\frac{(-1)^n}{4^n}C_n(x),\quad (n\geq 0). \tag{2.1}$$

For $d\in\mathbb{N}$ with $d\equiv 1 \pmod 2$, we have

$$\int_{\mathbb{Z}_p}(1-4t)^{\frac{x+d}{2}}d\mu_{-1}(x)+\int_{\mathbb{Z}_p}(1-4t)^{\frac{x}{2}}d\mu_{-1}(x) \tag{2.2}$$

$$= 2\sum_{i=0}^{d-1}(1-4t)^{\frac{i}{2}}(-1)^i$$
$$= \sum_{n=0}^{\infty}\left(2\sum_{i=0}^{d-1}\binom{\frac{i}{2}}{n}(-4)^n(-1)^i\right)t^n.$$

Thus, by (1.9) and (2.2), we get

$$\sum_{n=0}^{\infty}(C_n(d)+C_n)\,t^n = \sum_{n=0}^{\infty}\left(2^{2n+1}\sum_{i=0}^{d-1}\binom{\frac{i}{2}}{n}(-1)^{n-i}\right)t^n. \tag{2.3}$$

Therefore, by (2.3), we obtain the following theorem.

Theorem 1. *For $n \geq 0$, $d \in \mathbb{N}$ with $d \equiv 1 \pmod 2$, we have*

$$C_n(d)+C_n = 2^{2n+1}\sum_{i=0}^{d-1}\binom{\frac{i}{2}}{n}(-1)^{n-i}.$$

Now, we observe that, for any $d \in \mathbb{N}$,

$$\int_{\mathbb{Z}_p}(1-4t)^{\frac{x+d}{2}}\,d\mu_{-1}(x) + \int_{\mathbb{Z}_p}(1-4t)^{\frac{x}{2}}\,d\mu_{-1}(x)$$
$$= \frac{2}{\int_{\mathbb{Z}_p}(1-4t)^{\frac{d}{2}x}\,d\mu_{-1}(x)}\int_{\mathbb{Z}_p}(1-4t)^{\frac{x}{2}}\,d\mu_{-1}(x). \tag{2.4}$$

From (1.8), we have

$$\int_{\mathbb{Z}_p}(1-4t)^{\frac{d}{2}x}\,d\mu_{-1}(x) = \frac{2}{1+(1-4t)^{\frac{d}{2}}} = \sum_{n=0}^{\infty}C_{n,d}t^n. \tag{2.5}$$

Let $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$, $w_2 \equiv 1 \pmod 2$.
By using double fermionic p-adic invariant integral on $\mathbb{Z}_p$, we get

$$\frac{\int_{\mathbb{Z}_p}\int_{\mathbb{Z}_p}(1-4t)^{\frac{w_1x+w_2y}{2}}\,d\mu_{-1}(x)\,d\mu_{-1}(y)}{\int_{\mathbb{Z}_p}(1-4t)^{\frac{w_1w_2}{2}x}\,d\mu_{-1}(x)}$$
$$= \frac{2\left((1-4t)^{\frac{w_1w_2}{2}}+1\right)}{\left((1-4t)^{\frac{w_1}{2}}+1\right)\left((1-4t)^{\frac{w_2}{2}}+1\right)}. \tag{2.6}$$

Now, we also consider the following fermionic p-adic invariant integral on $\mathbb{Z}_p$ associated with w-Catalan polynomials.

$$I = \frac{\int_{\mathbb{Z}_p}\int_{\mathbb{Z}_p}(1-4t)^{\frac{w_1x_1+w_2x_2+w_1w_2x}{2}}\,d\mu_{-1}(x_1)\,d\mu_{-1}(x_2)}{\int_{\mathbb{Z}_p}(1-4t)^{\frac{w_1w_2}{2}x}\,d\mu_{-1}(x)}$$
$$= \frac{2(1-4t)^{\frac{w_1w_2}{2}x}\left((1-4t)^{\frac{w_1w_2}{2}}+1\right)}{\left((1-4t)^{\frac{w_1}{2}}+1\right)\left((1-4t)^{\frac{w_2}{2}}+1\right)}. \tag{2.7}$$

From (2.3), (2.4), we note that, $w_1, d \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$,

$$\frac{\int_{\mathbb{Z}_p} (1-4t)^{\frac{dx}{2}} d\mu_{-1}(x)}{\int_{\mathbb{Z}_p} (1-4t)^{\frac{dw_1 x}{2}} d\mu_{-1}(x)} = \sum_{i=0}^{w_1-1} (1-4t)^{\frac{di}{2}} (-1)^i \tag{2.8}$$

$$= \sum_{k=0}^{\infty} \left(\sum_{i=0}^{w_1-1} \binom{\frac{di}{2}}{k} (-1)^i (-4)^k \right) t^k$$

$$= \sum_{k=0}^{\infty} S_{k,d}(w_1-1)(-4)^k t^k.$$

By (2.7) and (2.8), we get

$$I = \left(\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1(x_1+w_2 x)}{2}} d\mu_{-1}(x_1) \right) \left(\frac{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_2 x_2}{2}} d\mu_{-1}(x_2)}{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 w_2}{2} x} d\mu_{-1}(x)} \right) \tag{2.9}$$

$$= \left(\sum_{l=0}^{\infty} C_{l,w_1}(w_2 x) t^l \right) \left(\sum_{k=0}^{\infty} S_{k,w_2}(w_1-1)(-4)^k t^k \right)$$

$$= \sum_{n=0}^{\infty} \left(\sum_{l=0}^{n} (-4)^{n-l} C_{l,w_1}(w_2 x) S_{n-l,w_2}(w_1-1) \right) t^n.$$

On the other hand,

$$I = \left(\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_2(x_2+w_1 x)}{2}} d\mu_{-1}(x_2) \right) \left(\frac{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 x_1}{2}} d\mu_{-1}(x_1)}{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 w_2}{2} x} d\mu_{-1}(x)} \right) \tag{2.10}$$

$$= \left(\sum_{l=0}^{\infty} C_{l,w_2}(w_1 x) t^l \right) \left(\sum_{k=0}^{\infty} S_{k,w_1}(w_2-1)(-4)^k t^k \right)$$

$$= \sum_{n=0}^{\infty} \left(\sum_{l=0}^{n} (-4)^{n-l} C_{l,w_2}(w_1 x) S_{n-l,w_1}(w_2-1) \right) t^n.$$

Therefore, by (2.9) and (2.10), we obtain the following theorem.

Theorem 2. *For $n \geq 0$, $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$ and $w_2 \equiv 1 \pmod 2$, we have*

$$\sum_{l=0}^{n} (-4)^{n-l} C_{l,w_1}(w_2 x) S_{n-l,w_2}(w_1-1) = \sum_{l=0}^{n} (-4)^{n-l} C_{l,w_2}(w_1 x) S_{n-l,w_1}(w_2-1).$$

Let $w_1 = 1$. Then we have

$$C_{n,1}(w_2 x) = \sum_{l=0}^{n} (-4)^{n-l} C_{l,w_2}(x) S_{n-l,1}(w_2-1). \tag{2.11}$$

Therefore, by (2.11), we obtain the following corollary.

Corollary 3. *For $w_2 \in \mathbb{N}$ with $w_2 \equiv 1 \pmod 2$, we have*

$$C_n(w_2 x) = \sum_{l=0}^{n} (-4)^{n-l} C_{l,w_2}(x) S_{n-l}(w_2 - 1).$$

When $x = 0$ in Theorem 2, we have the following corollary.

Corollary 4. *For $n \geq 0$, $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$, $w_2 \equiv 1 \pmod 2$, we have*

$$\sum_{l=0}^{n} (-4)^{n-l} C_{l,w_1} S_{n-l,w_2}(w_1 - 1) = \sum_{l=0}^{n} (-4)^{n-l} C_{l,w_2} S_{n-l,w_1}(w_2 - 1).$$

From (2.7), we have

(2.12)

$$\begin{aligned}
&I \\
&= \left((1-4t)^{\frac{w_1 w_2}{2} x} \int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 x_1}{2}} d\mu_{-1}(x_1) \right) \left(\frac{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_2 x_2}{2}} d\mu_{-1}(x_2)}{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 w_2}{2} x} d\mu_{-1}(x)} \right) \\
&= \left((1-4t)^{\frac{w_1 w_2}{2} x} \int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 x_1}{2}} d\mu_{-1}(x_1) \right) \left(\sum_{l=0}^{w_1-1} (-1)^l (1-4t)^{\frac{w_2}{2} l} \right) \\
&= \sum_{l=0}^{w_1-1} (-1)^l \int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1}{2}\left(x_1 + w_2 x + \frac{w_2}{w_1} l\right)} d\mu_{-1}(x_1) \\
&= \sum_{n=0}^{\infty} \left(\sum_{l=0}^{w_1-1} (-1)^l C_{n,w_1}\left(w_2 x + \frac{w_2}{w_1} l \right) \right) t^n.
\end{aligned}$$

On the other hand,

(2.13)

$$\begin{aligned}
&I \\
&= \left((1-4t)^{\frac{w_1 w_2}{2} x} \int_{\mathbb{Z}_p} (1-4t)^{\frac{w_2 x_2}{2}} d\mu_{-1}(x_2) \right) \left(\frac{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 x_1}{2}} d\mu_{-1}(x_1)}{\int_{\mathbb{Z}_p} (1-4t)^{\frac{w_1 w_2}{2} x} d\mu_{-1}(x)} \right) \\
&= \left((1-4t)^{\frac{w_1 w_2}{2} x} \int_{\mathbb{Z}_p} (1-4t)^{\frac{w_2 x_2}{2}} d\mu_{-1}(x_2) \right) \left(\sum_{l=0}^{w_2-1} (-1)^l (1-4t)^{\frac{w_1}{2} l} \right) \\
&= \sum_{l=0}^{w_2-1} (-1)^l \int_{\mathbb{Z}_p} (1-4t)^{\frac{w_2}{2}\left(x_2 + w_1 x + \frac{w_1}{w_2} l\right)} d\mu_{-1}(x_2) \\
&= \sum_{n=0}^{\infty} \left(\sum_{l=0}^{w_2-1} (-1)^l C_{n,w_2}\left(w_1 x + \frac{w_1}{w_2} l \right) \right) t^n.
\end{aligned}$$

Therefore, by (2.12) and (2.13), we obtian the following theorem.

Theorem 5. *For $n \geq 0$, $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$, $w_2 \equiv 1 \pmod 2$, we have*

$$\sum_{l=0}^{w_1-1} (-1)^l C_{n,w_1}\left(w_2 x + \frac{w_2}{w_1} l\right) = \sum_{l=0}^{w_2-1} (-1)^l C_{n,w_2}\left(w_1 x + \frac{w_1}{w_2} l\right).$$

Setting $w_2 = 1$ in Theorem 5, we get the multiplication– type formula for the Catalan polynomials as follows:

$$C_n(w_1 x) = \sum_{l=0}^{w_1-1} (-1)^l C_{n,w_1}\left(x + \frac{1}{w_1} l\right).$$

References

1. H. W. Gould, *Sums and convolved sums of Catalan numbers and their generating functions,* Indian J. Math. 46 (2004), no. 2-3, 137-160.
2. R. Hampel, *On the problem of Catalan,* (Polish) Prace Mat. 4 (1960), 11-19.
3. Y. He and C. Wang, *Recurrence formulae for Apostol-Bernoulli and Apostol-Euler polynomials* Adv. Difference Equ. 2012, 2012:209, 16 pp.
4. S. Hyyro, *On the Catalan problem,* (Finnish) Arkhimedes 1963 (1963), no. 1, 53-54.
5. K. Inkeri, *On Catalan's problem,* Acta Arith. 9 (1964), 285-290.
6. D. S. Kim and K. H. Park, *Identities of symmetry for Euler polynomials arising from quotients of fermionic integrals invariant under S_3*, J. Inequal. Appl. 2010, Art. ID **851521**, 16 pp.
7. T. Kim, *Symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials*, J. Difference Equ. Appl. **14** (2008), 1267-1277.
8. T. Kim, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$*. Russ. J. Math. Phys. 1616 (2009), no.**1**, 93-96.
9. T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. 9 (2002), no.**3**, 288-299.
10. T. Kim, *On p-adic interpolating function for q-Euler numbers and its derivatives*, J. Math. Anal. Appl. 339 (2008), no. **1**, 598-608.
11. A. Natucci, *Ricerche sistematiche intorno al "teorema di Catalan",* (Italian) Giorn. Mat. Battaglini (5) 2(82) (1954), 297-300.
12. H. Ozden, I. Naci Cangul and Y. Simsek,*Multivariate interpolation functions of higher-order q-Euler numbers and their applications*, Abstract and Applied Analysis, Article Number: **390857** Published: 2008.
13. H. Ozden and Y. Simsek, *A new extension of q-Euler numbers and polynomials related to their interpolation functions*, Applied Mathematics Letters, Volume **21**, Issue 9, September 2008, Pages 934-939.
14. R. Rangarajan and P. Shashikala, *A pair of classical orthogonal polynomials connected to Catalan numbers,* Adv. Stud. Contemp. Math. (Kyungshang) 23 (2013), no. 2, 323-335.
15. A. Rotkiewicz, *Sur le probléme de Catalan,* (French) Elem. Math. 15 (1960), 121-124.
16. A. D. Sands, *On generalised Catalan numbers,* Discrete Math. 21 (1978), no 2. 219-221.

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail address: tkkim@kw.ac.kr

Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
E-mail address: dskim@sogang.ac.kr

Department of Applied Mathematics, Pukyong National University, Busan, Republic of Korea
E-mail address: seo2011@pknu.ac.kr